



Do any of the following apply to your patient?

Patient history

- · Cancer diagnosed under age 50
- Cancer diagnosed at any age in the context of significant family history (see right)
- · Bilateral or multiple primary cancers
- Rare cancer^a
- Ashkenazi Jewish ancestry^b

Family history


- ≥2 relatives with related cancer,° with 1 diagnosed under age 50
- ≥3 relatives with related cancer^c on the same side of the family
- Relative with a known familial mutation
- Ashkenazi Jewish ancestry

^a Male breast cancer, metastatic prostate cancer, ovarian cancer, pancreatic cancer, colorectal or uterine cancer with abnormal microsatellite instability/immunohistochemistry (MSI/IHC), pheochromocytoma, paraganglioma, 10 or more gastrointestinal polyps

^bIncreased risk for specific *BRCA1* and *BRCA2* mutations

[°] For a complete list of related cancer types, see the Hereditary Cancer Reference Guide

Choosing the right test: a quick guide to help with collaborative, informed decision-making



For assistance, contact Genomic Client Services at 1.866.GENE.INFO (1.866.436.3463) or QuestHereditaryCancer.com

The information presented is not intended to be a complete source for cancer risk in any patient. The patient's physician must determine the patient's risk based upon the patient's clinical assessment, history, and the physician's training and experience.

QuestDiagnostics.com

Quest, Quest Diagnostics, any associated logos, and all associated Quest Diagnostics registered or unregistered trademarks are the property of Quest Diagnostics. All third-party marks—® and TM—are the property of their respective owners. © 2023 Quest Diagnostics Incorporated. All rights reserved. MI7552 3/2023